

## “ROLE OF PHYSIOTHERAPY FOR COVID – 19 PATIENTS”

*Dr. Shaima Parveen Quadri*

*Assistant Professor, Department of Physiotherapy, Jayoti Vidyapeeth Women's University, Jaipur*

### Abstract

Coronavirus disease 2019 / COVID-19 a infectious disease is initially found in Asia (Hubei ,China). Later on it called as Middle East respiratory syndrome corona virus, or MERS - CoV and also known as Severe Acute Respiratory Syndrome (SARS). COVID-19 disease spreads universally. plan creators, researchers, health care management, authority, intensive care unit (ICU) specialists have to reinforcement for increased in volume of significant for COVID-19.

Physiotherapists and other healthcare proffessinals are involved with care and treatment of COVID-19 patients and show an important part in expectant treatment and care , functional mobility, posture. Objective of the study is discover current research authentication for the PT Rx for COVID-19.

**Keywords :-** Physiotherapy [PT] , Physiotherapy treatment [PT Rx], COVID-19, SARS

### Introduction

Corona virus disease is severe acute respiratory syndrome coronavirus-2, (SARSCOV-2), which is appeared in 2019 it is a infectious disease. It is different from previous respiratory virus in a manner that it seems to have approximately 2-10 days human to human transmission before a person is becoming symptomatic<sup>1</sup>. virus is transfer from one person to another by respiratory emission. Respiratory droplet transmission can occur when a *person is in* close contact ... *infected person* who has respiratory symptoms (e.g. *coughing* or *sneezing*) possible modes of transmission for SARS-CoV-2, including contact, droplet, airborne, fomite, fecal-oral, bloodborne, mother-to-child, and animal-to-human transmission. Possible modes of transmission for SARS-CoV-2, including contact, droplet, airborne, fomite, fecal-oral, bloodborne, mother-to-child, and animal-to-human transmission<sup>2</sup>. COVID-19 involved symptoms like fever, body ache, dry cough, fatigue, chills, headache, sore throat, loss of appetite, and loss of smell and in severe patients may involved more severe symptoms like high fever, severe cough, and shortness of breath, which often indicates pneumonia. COVID-19 individuals may have influenza-like, respiratory tract infection symptoms such as pyrexia-89%, coughing-68%, extreme tiredness-38%, increased mucus-34% and shortness of breath (SOB)-19%<sup>3</sup>. This new coronavirus was linked to a wet seafood market, recognized as etiologic agent who is presently named as SARS-CoV-2<sup>4,5</sup>. According to recent studies the corona virus has spreading at the time of writing this total number of corona virus cases 7,307,097, deaths held 111,311 and number recovered are 6,383,441 occurs in the INDIA. The Global Surveillance Interim guidance developed by WHO<sup>6</sup>. (1) Restricting large gatherings has helped in limiting the spread of the virus. As some countries begin to allow gatherings again, it has never been more important to remain vigilant. For organizers of mass gatherings, the key planning recommendations from WHO outline the necessary precautions and safety measures when hosting large crowds of people. (2) the person is required to admit in hospital if person have symptoms like fever, cough, sneez etc. or the person may have visited china in past days. (3) If the person have any respiratory syndrome from past 14 days :

(i) Contacted a positive or possible COVID-19 infected patient or (ii). Served or appeared in medical canter where cases with positive or possible COVID-19 acute respiratory illness individuals were being treated. A individual with acute respiratory illness (ARI) having pyrexia with coughing & SOB along with one of the succeeding: (i) Travel history to China within 14 days preceding to symptoms. (ii) A close bodily contact within 14 days with a COVID-19 positive patient. The novel COVID-19 guidelines document<sup>8</sup> the person should admitted in the isolated ward for the treatment with in the 14 days as symptoms may persist. health-care workers, physical-therapists, especially respiratory therapists, are also playing an important role in managing and caring novel COVID-19 patients. They are involved in conservative care, posture correction, mobilization and while training to wean of from the weaning from invasive mechanical ventilator support<sup>7</sup>. Physiotherapist is a key element of the madical team of active hospital services and intensive care unit. Physiotherapy may be useful in the treatment of respiration, in the treatment of COVID-19, to proven work to prevent or delay critical care. Therefore, our aim is to provide worldwide evidence of physiotherapists involvement in managing COVID-19 infected patients.

### Purpose

This document has been prepared to provide information to physiotherapists and acute care

Health care facilities about the potential role of physiotherapy in the management of hospital admitted patients with confirmed and/or suspected COVID-19. COVID-19 is a disease caused by a new coronavirus, primarily impacting the respiratory system. Symptoms of COVID-19 can range from mild illness to pneumonia. Some people will have mild symptoms and recover easily, while others may develop respiratory failure and/or become critically ill and require admission to ICU. Physical therapy might be applicable for COVID-19 patients presenting with profuse airway discharges which patients cannot clear individually. Patients having associated diseases (like neuromuscular disease, respiratory diseases, lung fibrosis etc.) leading to increased secretion or weak cough may also be benefited from physical therapy. Physical therapists working in ICU can help in clearing airways for patients who are ventilated and provide assistance in placing them in proper position<sup>2</sup>.

## **Materials and Methods**

This is by looking in the PubMed, Scopus and Pedro databases to neglect the physiotherapist's work in serious reflection units in the management of non-invasive ventilation (NIV) without neglecting the risk of contamination for health experts.

### **Friendly approach**

A global specialist in cardiorespiratory physiotherapy met to quickly prepare clinical recommendations for the management of COVID-19 physiotherapy. The Creator group met for the first time on June 10, 2020 to investigate the imperative for the global alignment of physical therapy for the purpose of acute consideration according to COVID-19. There have been immediately made efforts to create an explicit guide for physical therapists in acute viewing situations. The capacity includes acute and intensive hospital physiotherapy (all), rehabilitation mediation in the intensive care unit (all), organization of physical therapy

Web search and individual records established recently established rules for the management of COVID-19 for primarily sick patients from universal offices ( e.g., World Health Organization), social regulations responsible for physiotherapy until June 10, 2020. These have been used to provide rules advice on transaction proposals on the main evaluation of the introductory meeting<sup>8</sup>.

### **Integration Criteria**

Patient who takes physiotherapy

- Infective exacerbation of COPD with acute deterioration
- Controlled asthma with evidence of infection, retained secretions, increased work of breathing
- Infective exacerbation of bronchiectasis or cystic fibrosis with difficulty managing retained secretions
- Significant consolidation with compromised respiratory status
- Atelectasis causing respiratory insufficiency
- Resolving and productive pneumonia with ineffective cough
- Recent estuation with associated retention of sputum and deteriorating ABGs/SaO<sub>2</sub> or significant risk of deterioration of respiratory status

## **Result**

### **PART 1: Orientation to physiotherapy considerations, detection and management**

#### **Orientation to physiotherapy**

Follow your hospital's rules and regulations regarding referral to physical therapy for patients who have been admitted to physical therapy in an acute setting<sup>9</sup>.

#### **Detection and disposition of reported patients**

1. The physiotherapist must perform an in-depth assessment of cases using the documentation framework.
2. Patients should be classified according to "Management of patients in physiotherapy for acute care COVID 19,
3. Patients who are approved for the airway without indication should maintain routine physical therapy care in accordance with standard contamination control precautions<sup>10</sup>.
4. Patients who have had respiratory side effects but who have doubts about COVID-19 (low isolation) should be classified as "suspected COVID 19" until two tests are reported that determine that the tests are negative. .
5. Patients who have been approved for other reasons (heart, injury, orthopedics or neurology) and who have indications for the respiratory tract must be classified as "suspected COVID 19" until two test determinations are reported in each case. which are negative.
6. Patients with respiratory diseases (e.g., shortness of breath, hacking, fever, sputum formation) must always be classified as "suspicious COVID 19", in any case two provisions for negative laboratory test reports.

#### **Considerations on the management of physiotherapy**

All suspected or confirmed cases must be reported to the director / supervisor through the group leader with regular updates so that they can be recorded very well on a COVID physiotherapy tracker / database accessible to all physiotherapy managers and the head of the Department<sup>11</sup>.

Suspicious or positive cases of COVID-19 should be assigned to the "COVID-19 Physiotherapy Group" for review. This will help limit / prevent the spread of contamination by making ideal considerations<sup>12</sup>.

## Part 2: Physiotherapy management categories

### TYPE A: ventilated, calm / incapacitated patients

It contains patients who generally do not feel well, who are calm, unable to act and possibly bowing. Patients with gadgets ECMO (extracorporeal membrane oxygenation): the understanding of ECMO devices depends on their level of sedation and loss of movement control.

The main goals of promoting physical therapy at this stage are

- Limit the confusion of lying on your back.
- Promote oxygen supply.

1. Physical therapy management may include (but not be limited to) distance-from-motion (ROM) activity and a recovery situation. (Trials are limited on the use of a wide range of exercise practices in anticipation of muscle weakness ICU.)
2. Physiotherapists should limit presentation to these patients to avoid unnecessary use of PPE and the risk of cross-contamination.
3. Collaborate with MDT to implement common restoration and insurance methods to limit or stop the presentation.
4. The decision for physiotherapy treatment should be based on the patient's ability to rehabilitate himself and consideration of the risks in relation to the benefits.
5. Regular correspondence with MD T relating to the physiotherapy management.

### TYPE B: patients evacuated with negligible ventilation / sedation

Develop an individual treatment plan based on the patient's level of knowledge, participation and hemodynamic status.

2. The main objectives of supporting physiotherapy mediations in this phase are:
3. Prevent tangles from lying down
4. Promote oxygen supply.
5. Improve freedom of supply

Physical therapy management can include drug implementation, ROM activities and dynamic activation<sup>13</sup>. Limiting the aerosol production process and physiotherapists must assess the risk versus the benefit to start these processes. In case of essential and extraordinary precautionary measures should be taken followed by these procedures.

### TYPE C: Non-precisely ventilated patients

1. The ESE patients classifications isolated based on a conscious level, and practical freedom.
2. The basic goals of physical therapy intercession are
  - Reduce craftsmanship through relaxation.
  - Improve the lung limit.
  - Promote oxygen supply.
  - Improve the useful limit.

### TYPE C.1: unconscious and out of service patients

- The intercession of physical therapy can integrate latent ROM activities and a recovery situation.
- The movement and recurrence of physical therapy treatment depend on the patient's rehabilitation potential.
- Regular recovery from MDT if the patient adheres to the practice program.

### TYPE C.2: conscious, dynamic and ward patients<sup>14</sup>

The physical therapy mediations may include ROM exercises, dynamic strengthening exercises, dynamic assembly and activities to improve coordination and balance.

- Develop a plan individual treatment depending on the oxygen dependence of the patient, the muscle performance and functional freedom, the limit functional advanced and advanced autonomy.
- Mobilization should be considered a flight production technique because it can cause piracy or expulsion. The most extreme warning must be followed when implementing these strategies.
- Special portability allows you to stay in the isolation area for use by patients with COVID 19 in particular.
- Make sure you have the necessary resources (by hand - at work as a device) before starting the activation.

- The necessary mobility aids must be labeled and left in the patient's room, or cleaned and disinfected, if they are to be reused for different patients.
- Avoid the use of specific large devices as reasonably expected or disinfect them carefully every time they are used in patients with COVID 19.
- Regular recovery from MDT if the patient adheres to the practice program.
- Strict adherence to contamination control procedures and precautions during preparation.

#### **TYPE C.3: conscious, dynamic and free patients<sup>15</sup>**

- Physical therapy intercession can include ROM exercises, dynamic walking exercises and breathing.
- Develop an individual training program that depends on the patient's endurance.
- Restrict or limit the presentation by instructing the patient in a protected training program and asking him or her to freely continue the training program.
- Regular monitoring of the MDT to ensure patient compliance with the training program and changes based on FITT rules (repetition, performance, duration and type).

#### **PART 3: Tips for best therapeutic practices.**

##### **Respiratory mediation advice<sup>16,17</sup>.**

- Intercession of the respiratory tract as postural infiltration, respiratory exercise strategies and methods of emissions released to the process of aerosol production.
- Avoid / limit this intercession and physiotherapists must weigh the risk with the benefits deriving from the initiation of these mediations.
- If breathing needs to be observed precautions in the air are. Before entering the patient's room, you need to use the personal protection equipment, such as items for the head, the veil breathing N95 suits cleaned using disposable, clothing uses disposable, eye protection (goggles or face shield), gloves and shoe covers.
- The N95 respirator should be at their ease at the point where the customer must adapt (eg a seal suction), to limit the amount of particles that bypass the channel through the holes between the skin of the customer and the seal of the breathing mask .
- The respirator must be worn (worn) and removed (removed) effectively and worn during introduction.
- For all personal defense equipment (PPE) must meet the pollution control requirements, according to respects the appropriate strategies for extraction and transportation are.
- During patient preparation, physiotherapists must follow the back method (if possible) and be  $\geq 2$  m away from the patient (if possible) to be outside the "impact zone" cut line.
- In patients with acute hypoxicity they can dyspnoea even in the presence of an organization of oxygen  $>$  with a 10-15 l / min bearing cap remaining. In this situation, the use of high flow nasal oxygen (HFNO) or continuous positive pressure (CPAP) or non-invasive ventilation (NIV) can be helpful during all physical therapy procedures.
- When preparing patients with HFNO / NIV, consider the possible natural distribution of vaporized particles of infection.
- The facial veil / deposition veil is ideal for the nasal cannula when a patient is assembled (facial cover with oxygen flow up to 5 l / min, revised cover up to 10 l / min of O<sub>2</sub> or Venturi veil up to 60 % FiO<sub>2</sub>) to distribute the limit accounts. A careful blanket covers the patient's face. If this veil is dirty, remove it according to the anti-pollution regulations and use a different cover.
- If the nasal cannula is the primary alternative, it should be fine in the nostrils and the cannula should be carefully covered. A similar standard is the material when the patient uses HFNO treatment.
- If the patient is ventilated noninvasively (CPAP / BIPAP), make sure that no air is released before starting a physical therapy session. In addition, it must have care to ensure that no separate allocation of circular machines.
- The VIN cover attached to a circulatory T-barrel can be used for patients with respirators to improve oxygen immersion during training.
- Limit or a strategic distance from the use of mechanical devices, such as pulse spirometers, PEEP, seaming, retaining the armor that these processes can theoretically produce in the air and the increase occurs during breathing.
- In ventilated patients with tracheotomy and precise, only if necessary recommended aspiration in closed circuit. Bronchial traction movements must be performed with severe signs.

Positioning to improve oxygenation: The patient is best positioned in a semi-sitting or sitting position while moving from the reclined position. Variations in parallel pressure ulcers, in semi-inclined or inclined positions can be very useful. The positions

should be resolved in fixed movements to reduce and the muscle of the jewels to relax to the ventilation / perfusion ratio to promote . Pads / pads can be used to achieve ideal (eliminated) rest positions.

#### **Advice on preparing mediations<sup>18</sup>**

- Mobilization should be seen as a vaporized production system because it causes piracy or expulsion. It also requires close contact of the physiotherapist with patients. In this sense, it should be a serious alarm are active in the realization of these methods are and precautions to be followed in the air.
- When assembling carefully ventilated patients, outrageous considerations must be made to maintain the ventilation circuit during activation.
- If the patient is not mechanically ventilated (CPAP / BIPAP), make sure that no air is released before starting a physical therapy session. We must also pay attention to maintaining the machine allocation circle.
- Patients not exactly ventilated Wear a veil during physical therapy sessions. If this veil gets dirty, it should be removed immediately in accordance with the disease control standards and covered with another cover. Make sure the appropriate resources (hand - in - the - job as a hardware, such as portability), before starting preparation. Avoid sharing material between patients.

#### **Advice on Infection control rehearses<sup>19</sup>**

Examine and follow approaches to control hospital contamination.

1. Remove, by way of example, any device / object close to home that contains earrings, watches, cables, mobile phones, acoustic signals and pens before entering the clinic area.
2. The reusable faculty defense equipment must be cleaned and disinfected before use.
3. In order to avoid cross-contamination, an immediate section of the area to be treated and locally defined documents are needed .
4. Physiotherapists who treat patients with COVID 19 are encouraged to wear clean coveralls or one-way coveralls that need to be cleaned daily before starting treatment.
5. Counsellors need to tie their hair and male specialists are asked to lose facial hair.
6. All physiotherapists must undergo an N95 fit test to determine the correct veil size.
7. All physiotherapists must complete the "use and remove" skill of the portable PSA and hand cleaning.

#### **Procedures should be applied during the acute phase with a blatant alarm<sup>20</sup>**

It is essential that physiotherapy procedures do not lead to weight gain during the manufacture of breathing. In COVID-19 patients, the primary goal of respiratory physiotherapy is to reduce the signs of shortness of breath and improve lung capacity, neutralizing the discomfort caused by respiratory illusions and immobilization, reduction of disability, improvement of personal satisfaction and reduction of the level of nervousness and discouragement of balance. After completing a physical therapy assessment, such as examining airway conditions, the physiotherapist must decide questions that are useful in the patient's treatment process based on clinical purposes. The point of administration of physical therapy at this stage of the infection movement may include:

- Reduce unnecessary respiratory activities (for example, with respiratory strategies and places that reduce wheezing, the oblique position, the course).
- clean the rest of the program,
- increase in the lung limit,
- improve the gas trade,
- gradually mobilize and expand the campaign,

General well-being acts, prevents the effects of immobilization and increases resistance to physical movements.

Postural situations to reduce shortness of breath Shortness of breath caused by cheating breathing can decrease and immobilize the patient's limit of activity. By placing the patient in a forward tilted position, we can rationalize the effects of the respiratory muscles and reduce the impression of dyspnea. It is recommended to prepare the patient out of bed if the persistent limit allows it.

#### **Respiratory system to reduce shortness of breath<sup>21</sup>:**

Breath control - this is supposed to be done. In the case of the patient's condition does not take such a position into account, act on a semi-reclining position (as above). It is important that the patient relaxes the inspiratory flying muscles, in particular those of the arms and neck, at this point he begins to breathe through the nose (warms and hydrates the air), at this point he plays a moderate relaxation and widens at the expiration with the establishment of adequate airways (lower thoracic breathing and stomach). Breathing should be calm (shallow, slow).

The conclusion must be made with the alleged relaxing lip of the lips. This method can be added for normal breathing while you rest and keep this movement in mind. It serves to slightly fix the lips while you exhale.

Physiotherapists work in a variety of environments and regardless of how the CPI can be the same for everyone and in all environments, the rest of the work will help reduce the need to do is in clinical centres, able to distinguish the activity of a physiotherapist in any context. In the basic idea (e.g. private offices, joint specialist studies or general practitioners), the highlight will be the rapid and visible identification and confirmation of cases. In taking care of society (e.g. at home), the supplement is to show patients and healthcare professionals. In serious reflection (for example, in the clinical centre), the complement will be in the organization of respiratory reactions.

## Discussion

The current epidemiological situation is an excellent test for any clinical expert. Now it is estimated that about 15 to 20 percent of patients with Covid-19 need to be hospitalized and 5-6 percent need a serious consideration for a period of time longer. As is common with this type of pandemic, the clinical staffs themselves are particularly contaminated and must take satisfactory measures to verify it.

Physiotherapists are an essential element in the multidisciplinary group of active hospital services and intensive care units. Early rehabilitation after the acute period of respiratory pain can limit the discomfort of stability in this way and promote rapid practical recovery. The expectation of this proposed good practice is to soften and improve the care physical therapy of patients with acute COVID and, at the same time, focus on the need to adhere to the undeniably well-established welfare rules<sup>1,19,20,21</sup>.

## Conclusion

The written survey showed the importance of physiotherapists in treating patients who are fundamentally critical in ICU. The ICU is a unique field in which physiotherapists are an essential part of the multidisciplinary group; the base camp offers various types of mediation, from physiotherapy in the thoracic phase, to the prevention and recovery of lack of strength. The primary goal of intensive care physiotherapy is the type of remote regeneration rather than temporary resistance, and physiotherapists do a lot of work to achieve this. It also seems essential to characterize teaching methods for healthcare professionals in order to receive adequate teaching, as well as for patients who require training on the use of equipment, if common rules can be established. Finally, due to the contamination of flight routes, the use of PSA should not be neglected, since in physiotherapy procedures, in which mist concentrates are generated, there is a high risk of transmission due to the contamination that is now Observe the coronavirus SARS 2 and in this way a risk is conceivable by the physiotherapists who use them.

## References

1. Shamsi S, Al-Shehri A, Khan S, Al Torairi N, Al Amoudi KO. Importance of Physiotherapy in COVID-19: A Recommendation. *Int J Rec Innov Med Clin Res*. 2020; 2(3):46-54.
2. Thomas P, Baldwin C, Bissett B, Boden I, Gosselink R, Granger CL, Hodgson C, Jones AY, Kho ME, Moses R, Ntoumenopoulos G. Physiotherapy management for COVID-19 in the acute hospital setting: clinical practice recommendations. *J Physiother*. 2020; 66(2):73-82.
3. Guan WJ, Ni ZY, Hu Y, Liang WH, Ou CQ, He JX, Liu L, Shan H, Lei CL, Hui DS, Du B. Clinical characteristics of coronavirus disease 2019 in China. *N Engl J Med*. 2020;30;382(18):1708-20.
4. Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, Zhang L, Fan G, Xu J, Gu X, Cheng Z. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *The Lancet*. 2020;15;395(10223):497-506.
5. Zhu N, Zhang D, Wang W, Li X, Yang B, Song J, Zhao X, Huang B, Shi W, Lu R, Niu P. A novel coronavirus from patients with pneumonia in China, 2019. *N Engl J Med*. 2020; 382:727– 33.
6. World Health Organization. Global Surveillance for human infection with novel coronavirus (2019-nCoV): Interim guidance v3. Geneva, Switzerland: World Health Organization; 2020. Available from: WHO/2019- nCoV/SurveillanceGuidance/ 2020.3.
7. Lazzeri M, Lanza A, Bellini R, Bellofiore A, Cecchetto S, Colombo A, D'Arosca F, Del Monaco C, Gaudellio G, Paneroni M, Privitera E. Respiratory physiotherapy in patients with COVID-19
8. Graziano Onder, Giovanni Rezza, and Silvio Brusaferro. Case-fatality rate and characteristics of patients dying in relation to COVID-19 in Italy. 2020;JAMA.
9. Michael Day. Covid-19: identifying and isolating asymptomatic people helped eliminate virus in Italian village. *The British Medical Journal*, 2020;3;68:165.
10. Chen N., Zhou M., Dong X, Qu J., Gong F., Han Y, et al. Epidemiological and clinical characteristics of 99 cases of new coronavirus pneumonia 2019 in Wuhan, China: a descriptive study. *The Lancet*. 2020; 395: 507-513.
11. 11 Kenji Mizumoto, Katsushi Kagaya, Alexander Zarebski, and Gerardo Chowell. Estimating the asymptomatic proportion of coronavirus disease 2019 (COVID-19) cases on board the Diamond Princess cruise ship, Yokohama, Japan, 2020; *Eurosurveillance*, 25(10), 2020
12. Zhou F., Yu T., Du R., Fan G., Liu Y, Liu Z. et al. Clinical development risk factors and mortality in adults hospitalized with COVID-19 in Wuhan, China: a retrospective cohort study. *The Lancet*. 2020;11.
13. Xie J, Tong Z, Guan X, Du B, Qiu H, Slutsky AS. Crisis in intensive care and some recommendations during the COVID-19 epidemic in China. *Intensive care med*. 2020;2.
14. Lesley Gibson and David Rush. Novel Coronavirus in Cape Town informal settlements: Feasibility of using informal dwelling outlines to identify high risk areas for COVID-19 transmission from a social distancing perspective. *JMIR Public Health and Surveillance*, 2020;6:2.
15. 15 Anne-Lise Sibony. The UK COVID-19 response: A behavioural irony? *European Journal of Risk Regulation*, 2020;1–11.
16. 16 Brouwers MC, Kho ME, Browman GP, Burgers JS, Cluzeau F., Feder G. et al. AGREE II development, part 1: performance, usefulness and improvement opportunities. *CMAJ*. 2010; 182 (10): 1045-105; 1052.
17. Joseph T Wu, Kathy Leung, and Gabriel M Leung. Nowcasting and forecasting the potential domestic and international spread of the 2019-nCoV outbreak originating in Wuhan, China: a modelling study. *The Lancet*, 2020;395(10225):689–697.
18. Schünemann HJ, Wiercioch W, Brozek J, Etzeandía-Ikobaltzeta I, Mustafa RA, Manja V, et al. GRADE Evidence to Decision (EtD) for the acceptance, adaptation and development of for health system and public health decisions:GRADE-ADOLOPMENT. *J ClinEpidemiol*. 2017; 81: 101-110.

19. .ShiZhao, QianyinLin, JinjunRan, SalihuSMusa, GuangpuYang, Weiming Wang, YijunLou, Daozhou Gao, Lin Yang, DaihaiHe, etal. Preliminary estimation of the basic reproduction number of novel coronavirus (2019-nCoV) in China, from 2019 to 2020: A data-driven analysis in the early phase of the outbreak. *International Journal of Infectious Diseases*, 2020;92: 214–217.
20. 20 Moberg J., Oxman AD, Rosenbaum S., Schünemann H. J., Guyatt G., Flottorp S., et al. The GRDE evidence to the decision framework ( ETD ) for the health system and public health decisions. *Syst. Res. Directive* 2018; 4:45.
21. Clinical skills development service, online physiotherapy and intensive care training. Available in <https://central.csd.s.qld.edu.au/central/courses/108>. [Access to March 21, 2020].
22. World Health Organization. Prevention and control of healthcare infections when Covid-19 is suspected: Intermediate guidelines. 2020. [Access to March 21, 2020].